

Friday 27 May 2022 – Morning

GCSE (9–1) Chemistry B (Twenty First Century Science)

J258/03 Breadth in Chemistry (Higher Tier)

Time allowed: 1 hour 45 minutes



You	must	have:

- a ruler (cm/mm)
- the Data Sheet for GCSE (9–1) Chemistry B (inside this document)

You can use:

• an HB pencil

• a scientific or graphical calculator



Please write clearly in black ink. Do not write in the barcodes.										
Centre number	Centre number Candidate number									
First name(s)										
Last name										

INSTRUCTIONS

- Use black ink. You can use an HB pencil, but only for graphs and diagrams.
- Write your answer to each question in the space provided. If you need extra space use the lined pages at the end of this booklet. The question numbers must be clearly shown.
- Answer all the questions.
- Where appropriate, your answer should be supported with working. Marks might be given for using a correct method, even if your answer is wrong.

INFORMATION

- The total mark for this paper is **90**.
- The marks for each question are shown in brackets [].
- This document has 28 pages.

ADVICE

• Read each question carefully before you start your answer.

Answer all the questions.

1 Mia investigates the rate of reaction when zinc reacts with dilute sulfuric acid. She adds zinc pieces to dilute sulfuric acid at room temperature.

Fig. 1.1 shows the apparatus she uses:



Fig. 1.1

(a) Complete the symbol equation for the reaction.

Include a state symbol.

$$Zn(s) + H_2SO_4(aq) \rightarrow ZnSO_4(aq) + \dots$$

[2]

(b) Mia measures the volume of gas in the gas syringe every two minutes.

Fig. 1.2 shows a graph of her results:



- 3
- (i) Calculate the rate of reaction during the first minute, using Fig. 1.2.
 Give your answer in cm³/s.

 (c) Mia repeats the experiment at the same temperature. She adds a few drops of blue copper sulfate.

Her results show that:

- more gas is produced in the first minute, than in the first experiment.
- the blue colour changes to colourless during the reaction.

Mia makes a statement about the results:



How do the results support Mia's statement?

Explain your answer.

 	 [3]

(d) Mia repeats the experiment at a higher temperature.

Which statements explain why the reaction is faster at a higher temperature?

Tick (✓) **two** boxes.

The particles move faster.

There are more frequent collisions.

The yield is higher at a higher temperature.

The particles are closer together.

The zinc breaks down into smaller pieces.

Γ

[2]

2 The table shows some properties of metals, polymers and clay ceramics:

Type of material	Effect of force on material	Electrical conductivity	Hardness	
Metals	malleable	good	hard	
Polymers	flexible	poor	soft	
Clay ceramics	snaps			

(a) Complete the table by adding the two missing properties of clay ceramics.

[2]

(b) Layla has three different water jugs. The jugs are made from aluminium, poly(ethene) and pottery (clay ceramic).

Draw lines to connect each material with its correct property.

Material

Aluminium

Poly(ethene)

Pottery

Property of jug

Softens easily when heated.

Goes out of shape if dropped.

Breaks if dropped.

Rusts quickly.

[3]

(c) Poly(ethene) is made from ethene.

Complete the equation by drawing the structure of the repeating unit of poly(ethene).



Ethene

Poly(ethene)

[1]

(d) Genes are made from the natural polymer DNA.

Which monomers make DNA?

Tick (✓) one box.

Cellulose

Nucleotides

Sugars

Unsaturated hydrocarbons

[1]

- 3 The Earth's early atmosphere contained mostly carbon dioxide and water vapour.
 - (a) As the Earth cooled, the oceans formed. Water vapour condensed from gas to liquid.

The particle model can explain properties of gases and liquids.

Which statement is correct?

Tick (✓) **one** box.

Liquids are difficult to squash because the particles are tightly packed.

Gases have pressure because gas particles rarely collide with each other.

Liquids expand when heated because atoms expand when heated.

Gases are easy to squash because gas atoms get smaller when squashed.

[1]

[1]

(b) The particle model assumes molecules of water and carbon dioxide are inelastic spheres. This is a limitation of the particle model.

Water changes state at 100 °C. Carbon dioxide changes state at -79 °C.

Which statement explains another limitation of the particle model in relation to changes of state?

Tick (✓) **one** box.

In real life, all molecules are the same size.

In real life, the molecules of water and carbon dioxide attract each other.

In real life, molecules become smaller during evaporation.

In real life, evaporation happens when bonds are broken in molecules.

(c) Gradually, plants began to grow on Earth. Explain how this affected the composition of the atmosphere.

.....[2]

(d) The graph shows how the percentage of carbon dioxide in the atmosphere has changed over time:



Ling and Ali discuss the graph:



Ling	
Ali	
	[2]

- 4 When hydrogen burns, energy is transferred to the surroundings.
 - (a) What word describes a reaction that releases energy to the surroundings?
 -[1]
 - (b) The equation for the reaction is:

 $2H_2(g) + O_2(g) \rightarrow 2H_2O(I)$

Complete **Fig. 4.1** to show the reaction profile for this reaction. Include on **Fig. 4.1**:

- the formulae of the reactants and products
- a label for the activation energy.



Progress of the reaction

- Fig. 4.1
- (c) Which statement describes the activation energy?

Tick (✓) one box.

The energy absorbed by the reaction.

The energy given out by the reaction.

The energy needed for a reaction to occur.

The temperature needed to start the reaction.

[1]

[2]

11

(d) Fig. 4.2 shows what happens when 2 moles of hydrogen burn:



Fig. 4.2

Calculate the energy given out when 2 moles of hydrogen burn.

Use data from the table.

Bond	Energy (kJ/mol)
O–H	464
O=O	498
H–H	436

Bonds broken =	kJ/mol
Bonds made =	kJ/mol
Energy given out =	kJ/mol [3]

- **5** Iodine is a halogen. It is an element in Group 17 (7) of the Periodic Table.
 - (a) Which property shows iodine is a non-metal?

Tick (✔) **one** box.

It is a shiny grey solid.	
It does not conduct electricity.	
It is unreactive.	
Its density is higher than water.	

(b) 1.00 cm^3 of iodine has a mass of 4.93 g.

Calculate the volume of 1.00 mol of ${\rm I_2}.$

Use the equation: number of moles =

mass of substance (g) relative formula mass (g)

Give your answer to 1 decimal place.

Volume = cm³ [3]

[1]

(c) Sodium iodide solution reacts with chlorine. This is a symbol equation for the reaction:

2NaI(aq) + $Cl_2(aq) \rightarrow 2NaCl(aq) + I_2(aq)$

(i) What would you see when this reaction happens?

-[1]
- (ii) What information does this reaction show about the reactivity of the halogens?

......[1]

.....

(d) An isotope of iodine has a mass number of 127.

How many neutrons and electrons does an atom of this isotope have?

Use the Data Sheet.

Number of neutrons =

Number of electrons =

[2]

(e) Phosphorus reacts with iodine to form a compound.

The compound contains 7.5% by mass of phosphorus. The rest of the compound is iodine.

Determine the formula of the compound.

Use the equation: number of moles = $\frac{\text{mass of substance }(g)}{\text{relative formula mass }(g)}$

Formula =[3]

- 6 In industry, crude oil fractions can be cracked to make alkenes.
 - (a) Which statement explains why crude oil fractions are cracked?

Tick (✓) one box.

Alkenes dissolve in water. Cracking separates the crude oil fractions. Crude oil is a mixture of alkenes.

To make useful short chain molecules.

(b) Bromine reacts with ethene.

Fig. 6.1 shows the fully displayed formula of ethene:

Н Н

Fig. 6.1

Draw the fully displayed formula of the product made when bromine reacts with ethene.

[1]

(c) (i) The reactions of organic compounds depend on their functional groups.

Put a (ring) around the functional groups present in ethene and ethanol in Fig. 6.2.



Fig. 6.2

[2]

(d) Ethanol boils at 78 °C. Ethene boils at –104 °C.
Which statement explains why the boiling points are different?
Tick (✓) one box.
An ethanol molecule has more carbon atoms than an ethene molecule.
Ethanol is soluble in water but ethene is not.
The covalent bonds in ethanol are stronger than those in ethene.
The intermolecular forces in ethanol are stronger than those in ethene.

[1]

[1]

7 (a) Iron ore contains iron oxide. Iron can be made from iron ore. This is a symbol equation for the reaction:

 $\mathrm{Fe_2O_3}$ + 3C \rightarrow 2Fe + 3CO

(i) Complete the half equation to show what happens to iron in this reaction.

 Fe^{3+} + $e^- \rightarrow$

- (ii) Explain, in terms of electrons, why Fe³⁺ is reduced in the half equation in (a)(i).
 -[1]

[1]

(b) Calculate the percentage by mass of iron present in iron oxide, Fe_2O_3 .

Give your answer to **3** significant figures.

	Percentage of Fe = % [3]
(c)	Iron can be made into steel. The main disadvantage of steel is that it rusts.
	Explain one method of preventing steel from rusting.
	[2]
(d)	Used steel is collected and recycled. Recycling used steel has advantages and disadvantages.
	Suggest one advantage and one disadvantage of recycling used steel.
	Advantage
	Disadvantage
	[2]

8 Magnesium reacts with iron sulfide. This is a symbol equation for the reaction:

 $Mg + FeS \rightarrow Fe + MgS$

(a) What information does this reaction give about the reactivity of magnesium?

......[1]

(b) Magnesium sulfide, MgS is an ionic compound which contains the ions Mg²⁺ and S²⁻. Explain why magnesium sulfide has a high melting point.

(c) Complete Fig. 8.1 to show the 'dot and cross' diagrams for an Mg²⁺ ion and an S²⁻ ion.
 Show all the electrons.



Fig. 8.1

[2]

- **9** Alex adds the **same** volume of dilute sulfuric acid to different samples of zinc carbonate. The reaction fizzes.
 - (a) Name the gas that causes the fizzing.

......[1]

(b) Table 9.1 shows Alex's results:

Experiment	Mass of zinc carbonate (g)	Concentration of acid (mol/dm ³)	Type of zinc carbonate	Time to stop fizzing (minutes)	Relative average rates
1	2.0	1.0	lumps	10	1
2	2.0	1.0	powder	4	

Table 9.1

(i) Explain why the rates of **Experiment 1** and **Experiment 2** are different in **Table 9.1**.

Use ideas about particles in your answer.

(ii) Relative average rate is the number of times faster one reaction is compared to another.Calculate the relative average rate of Experiment 2, compared to Experiment 1.

(c) Alex does two more experiments, **Experiment 3** and **Experiment 4**.

The results are shown in **Table 9.2**.

Experiment	Mass of zinc carbonate (g)	Concentration of acid (mol/dm ³)	Type of zinc carbonate	Time to stop fizzing (minutes)	Relative average rates
1	2.0	1.0	lumps	10	1
2	2.0	1.0	powder	4	
3	4.0	1.0	lumps		
4	2.0	2.0	lumps		

Table 9.2

Complete **Table 9.2** by predicting the time taken for the reactions to stop fizzing. [2]

10 Nina is given some diluted drain cleaner called 'Drainclear'. 'Drainclear' contains sodium hydroxide.

Nina titrates 25.0 cm³ of diluted 'Drainclear' with dilute hydrochloric acid and an indicator.

(a) Nina uses a burette to measure out the dilute hydrochloric acid in the titration.

Nina wants to minimise errors in her method.

Describe **one** thing she should do when taking burette readings.

[1]

(b) The table shows Nina's results:

	Rough	Titration 1	Titration 2	Titration 3
2nd reading (cm ³)	14.05	20.55	10.60	22.05
1st reading (cm ³)	3.00	10.05	0.05	11.60
Volume (cm ³)	11.05	10.50	10.55	10.45
Mean (cm ³)		10.50		

Explain why Nina's results are repeatable but not reproducible.

......[2]

(c) In her titration, Nina used 25.0 cm³ of diluted 'Drainclear' with 0.20 mol/dm³ hydrochloric acid. A symbol equation for the reaction is:

 $NaOH(aq) + HCl(aq) \rightarrow NaCl(aq) + H_2O(I)$

Calculate the concentration of sodium hydroxide, NaOH, in the diluted 'Drainclear'.

Use the equation: concentration (mol/dm³) = $\frac{\text{number of moles of solute}}{\text{volume (dm³)}}$

<u>_</u>

Use Nina's mean result of 10.50 cm³.

Concentration of sodium hydroxide = mol/dm³ [3]

(d) Nina says:

'I would have preferred my titration result to be larger than 10.50 cm³. This would reduce the percentage uncertainty in my titration result.'

Explain how Nina could get a larger titration result without changing her apparatus.

[2]

22

- **11** Aluminium and chlorine can be made by electrolysis.
 - (a) Aluminium is made by the electrolysis of molten aluminium oxide.

	(i)	At which electrode is aluminium formed?
		[1]
	(ii)	Give one reason for your answer to (a)(i) .
		[1]
(b)	Sor	ne elements are extracted using electrolysis.
	(i)	Explain why hydrogen, not sodium, is made at the negative electrode when sodium chloride solution is electrolysed.
		[1]
	(ii)	Write the half equation for the formation of hydrogen at the negative electrode.
		[2]

Describe what Kareem needs to do to measure the volumes of hydrogen gas and chlorine gas made.

You may add to the diagram to help explain your answer. The diagram is **not** complete.



 [2]

- **12** Eve has some copper sulfate crystals. The formula of copper sulfate is CuSO₄.
 - (a) Eve says, 'Copper sulfate is a mixture of several elements. It is not a pure substance.'

Explain why Eve is wrong.



END OF QUESTION PAPER

ADDITIONAL ANSWER SPACE

If additional space is required, you should use the following lined page(s). The question number(s) must be clearly shown in the margin(s).

•••••	

.....

.....

.....I.

.....

.....

.....I.

.....



Copyright Information

OCR is committed to seeking permission to reproduce all third-party content that it uses in its assessment materials. OCR has attempted to identify and contact all copyright holders whose work is used in this paper. To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced in the OCR Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download from our public website (www.ocr.org.uk) after the live examination series. If OCR has unwittingly failed to correctly acknowledge or clear any third-party content in this assessment material, OCR will be happy to correct its mistake at the earliest possible opportunity.

_____I

For queries or further information please contact The OCR Copyright Team, The Triangle Building, Shaftesbury Road, Cambridge CB2 8EA.

OCR is part of Cambridge University Press & Assessment, which is itself a department of the University of Cambridge.

.....